

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number
WO 01/00231 A2

(51) International Patent Classification⁷: **A61K 39/00**

(21) International Application Number: **PCT/EP00/05841**

(22) International Filing Date: **23 June 2000 (23.06.2000)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
9915204.3 **29 June 1999 (29.06.1999)** **GB**

(71) Applicant (*for all designated States except US*):
SMITHKLINE BEECHAM BIOLOGICALS S.A.
[BE/BE]; Rue de L'Institut 89, B-1330 Rixensart (BE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **COHEN, Joseph**
[US/BE]; SmithKline Beecham Biologicals S.A., Rue de
l'Institut 89, B-1330 Rixensart (BE). **GARCON, Nathalie**
[FR/BE]; SmithKline Beecham Biologicals S.A., Rue
de l'Institut 89, B-1330 Rixensart (BE). **VOSS, Gerald**
[DE/BE]; SmithKline Beecham Biologicals S.A., Rue de
l'Institut 89, B-1330 Rixensart (BE).

(74) Agent: **TYRRELL, Arthur, William, Russell**; SmithK-
line Beecham, Corporate Intellectual Property, Two New
Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— *Without international search report and to be republished
upon receipt of that report.*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: **VACCINES**

(57) Abstract: A vaccine formulation for the prevention or amelioration of plasmodium infection in humans is provided. The vaccine comprises a malaria antigen, especially a protein which comprises a portion of the CS protein of *P. falciparum* fused in frame via a linear linker to the N-terminal of HBsAg, and an immunostimulatory CpG oligonucleotide. Methods for making the vaccine formulation of the invention are described. Patients may also be treated by pre-administration of the CpG oligonucleotide prior to administration of the malaria antigen.

BEST AVAILABLE COPY

WO 01/00231 A2

VACCINES

The present invention relates to a novel vaccine formulations and their use in medicine, particularly in the prevention of malaria infections. In particular the present invention is concerned with a CpG oligonucleotide and a malarial antigen.

Malaria, is one of the world's major health problems with 2 to 4 million people dying from the disease each year. One of the most acute forms of the disease is caused by the protozoan parasite, Plasmodium falciparum which is responsible for most of the mortality attributable to Malaria.

The life cycle of P. falciparum is complex, requiring two hosts, man and mosquito for completion. The infection of man is initiated by the inoculation of sporozoites in the saliva of an infected mosquito. The sporozoites migrate to the liver and there infect hepatocytes where they differentiate, via the exoerythrocytic intracellular stage, into the merozoite stage which infects red blood cells (RBC) to initiate cyclical replication in the asexual blood stage. The cycle is completed by the differentiation of a number of merozoites in the RBC into sexual stage gametocytes which are ingested by the mosquito, where they develop through a series of stages in the midgut to produce sporozoites which migrate to the salivary gland.

The sporozoite stage of P. falciparum has been identified as a potential target of a malaria vaccine. The major surface protein of the sporozoite is known as circumsporozoite protein (CS Protein). This protein from strain 7G8 has been cloned, expressed and sequenced (Dame et al Science 225 (1984) p593). The protein from strain 7G8 is characterised by having a central immunodominant repeat region comprising a tetrapeptide Asn-Ala-Asn-Pro repeated 37 times but interspersed with four minor repeats Asn-Val-Asp-Pro. In other strains the number of major and minor repeats vary as well as their relative position. This central portion is flanked by an N and C terminal portion composed of non-repetitive amino acid sequences designated as the repeatless portion of the CS protein.

It has been shown that irradiated sporozoites can provide significant protection against experimental human malaria (Am. J. Trop. Med. Hyg. 24: 297-402, 1975). However, production difficulties makes the use of irradiated sporozoite impractical from the point of view of producing a vaccine.

5

Several groups have proposed subunit vaccines based on the circumsporozoite protein. Two of these vaccines have undergone clinical testing; one is a synthetic peptide, the other is a recombinant protein (Ballou et al Lancet: i 1277 (1987) and Herrington et al Nature 328:257 (1987).

10

These vaccines were successful in stimulating an anti-sporozoite response. Nonetheless, the magnitude of the response was disappointing, with some vaccinees not making a response at all. Furthermore, the absence of "boosting" of antibody levels on subsequent injections and results of in vitro lymphocyte proliferation assays suggested that T-cells of most of these volunteers did not recognise the immuno-dominant repeat. Nonetheless, one vaccinee in each study did not develop parasitemia.

15

The present invention provides a new, improved malaria vaccines which not only produces a humoral response, but also a cellular immune response. Preferably the antigen induces the production of neutralising antibodies against the immunodominant repeat. Most preferably, the antigen should also elicit effector T cell mediated immune responses of the CD4⁺ and CD8⁺ cytotoxic T lymphocyte (CTL) type and of the delayed type hypersensitivity type and also, preferably be able to induce T helper (TH) memory cells.

20

25

International patent application No. WO 93 / 10152 (SmithKline Beecham Biologicals s.a) provides a hybrid protein comprising substantially all the C-terminal portion of the CS protein, four or more tandem repeats of the immunodominant region, and the Surface antigen from Hepatitis B virus (HBsAg). Preferably the hybrid protein comprises a sequence which contains at least 160 amino acids which is substantially homologous to the C-terminal portion of the CS

30

protein. The CS protein may be devoid of the last 12 amino-acids from the C terminal.

In particular there is provided a protein which comprises a portion of the CS protein of *P. falciparum* substantially as corresponding to amino acids 210-398 of *P. falciparum* 7G8 fused in frame via a linear linker to the N-terminal of HBsAg. The linker may comprise a portion of preS2 from HBsAg.

A particularly preferred embodiment is the hybrid protein designated RTS (or RTS,S). This hybrid consists of:

- A methionine-residue, encoded by nucleotides 1059 to 1061, derived from the *Saccharomyces cerevisiae* TDH3 gene sequence. (Musti A.M. et al Gene 1983 25 133-143).
- Three amino acids, Met Ala Pro, derived from a nucleotide sequence (1062 to 1070) created by the cloning procedure used to construct the hybrid gene.
- A stretch of 189 amino acids, encoded by nucleotides 1071 to 1637 representing amino acids 210 to 398 of the circumsporozoite protein (CSP) of *Plasmodium falciparum* strain 7G8 (Dame et al supra).
- An amino acid (Arg) encoded by nucleotides 1638 to 1640, created by the cloning procedure used to construct the hybrid gene.
- Four amino acids, Pro Val Thr Asn, encoded by nucleotides 1641 to 1652, and representing the four carboxy terminal residues of the hepatitis B virus (adw serotype) preS2 protein (9).
- A stretch of 226 amino acids, encoded by nucleotides 1653 to 2330, and specifying the S protein of hepatitis B virus (adw serotype).

In an alternative embodiment there is provided a hybrid protein designated RTS* (or RTS*,S), which was generated using the CSP gene sequence from *P. falciparum* NF54 (Mol. Biochem Parasitol. 35 : 185-190, 1989) and comprises substantially all of the region 207 to 395 of the CS protein from *P falciparum* NF54.

5

In particular RTS* comprises:

10

- A Methionine, encoded by nucleotides 1059 to 1061, derived from the TDH3 gene sequence (see Musti et al, loc cit)..

15

- Three amino acids, Met Ala Pro, derived from a nucleotide sequence (1062 to 1070) created by the cloning procedure used to construct the hybrid gene.
- A stretch of 189 amino acids, encoded by nucleotides 1071 to 1637 representing amino acids 207 to 395 of the circumsporozoite protein (CSP) of *Plasmodium falciparum* strain NF54 (Mol.Biochem.Parasitol, 35:185-190, 1989).

20

- An amino acid (Gly) encoded by nucleotides 1638 to 1640, created by the cloning procedure used to construct the hybrid gene.
- Four amino acids, Pro Val Thr Asn, encoded by nucleotides 1641 to 1652, and representing the four carboxy terminal residues of the hepatitis B virus (adw serotype) preS2 protein (Nature 280:815-819, 1979).

25

- A stretch of 226 amino acids, encoded by nucleotides 1653 to 2330, and specifying the S protein of hepatitis B virus (adw serotype) (Nature 280:815-819, 1979)

30

International patent application no. WO 90/01496 describe an antigen known as Trap (or TRAP) from *P. falciparum*. An apparent homologue of Trap is described in WO92/11868 and relates to an antigen called SSP2 from *P. yeolii*.

International patent application WO 98/ 05355 describes, inter alia, a malaria vaccine based on a combination of Trap and RTS,S.

- 5 Immunomodulatory oligonucleotides contain unmethylated CpG dinucleotides ("CpG") and are known (WO 96/02555, EP 468520). CpG is an abbreviation for cytosine-guanosine dinucleotide motifs present in DNA. Historically, it was observed that the DNA fraction of BCG could exert an anti-tumour effect. In further studies, synthetic oligonucleotides derived from BCG gene sequences were
- 10 shown to be capable of inducing immunostimulatory effects (both in vitro and in vivo). The authors of these studies concluded that certain palindromic sequences, including a central CG motif, carried this activity. The central role of the CG motif in immunostimulation was later elucidated in a publication by Krieg, Nature 374, p546 1995. Detailed analysis has shown that the CG motif has to be in a certain
- 15 sequence context, and that such sequences are common in bacterial DNA but are rare in vertebrate DNA.

- It is currently believed that this evolutionary difference allows the vertebrate immune system to detect the presence of bacterial DNA (as occurring during an
- 20 infection) leading consequently to the stimulation of the immune system. The immunostimulatory sequence as defined by Krieg is:

- Purine Purine CG pyrimidine pyrimidine and where the CG motif is not methylated. In certain combinations of the six nucleotides a palindromic sequence is present.
- 25 Several of these motifs, either as repeats of one motif or a combination of different motifs, can be present in the same oligonucleotide. The presence of one or more of these immunostimulatory sequence containing oligonucleotides can activate various immune subsets, including natural killer cells (which produce interferon γ and have cytolytic activity) and macrophages (Wooldrige et al Vol 89 (no. 8), 1977).
- 30 Although other unmethylated CpG containing sequences not having this consensus sequence have now been shown to be immunomodulatory.

The present invention provides an improved vaccine formulation comprising a CpG oligonucleotide and a malaria antigen. In particular, RTS,S or RTS,S* or Trap or immunologically equivalent derivatives thereof.

- 5 Vaccine preparation is generally described in Vaccine Design - The subunit and adjuvant approach (Ed. Powell and Newman) Pharmaceutical Biotechnology Vol. 6 Plenum Press 1995. Encapsulation within liposomes is described by Fullerton, US Patent 4,235,877.
- 10 The preferred oligonucleotides preferably contain two or more CpG motifs separated by six or more nucleotides. The oligonucleotides of the present invention are typically deoxynucleotides. In a preferred embodiment the internucleotide in the oligonucleotide is phosphorodithioate, or more preferably a phosphorodithioate bond, although phosphodiester and other internucleotide bonds are within the scope
- 15 of the invention including oligonucleotides with mixed internucleotide linkages. The sequences preferably contain all phosphorodithioate modified internucleotide linkages. Preferred oligonucleotides have the following sequences:

Oligo (internal designation*)	5'-SEQUENCE-3'	CpG	Thio
WD1001	TCC ATG ACG TTC CTG ACG TT	+	+
WD1002	TCT CCC AGC GTG CGC CAT	+	+
WD1003	ACC GAT AAC GTT GCC GGT GAC G	+	-
WD1004	G*G*G GTC AAC GTT GAG* G*G*G* G*G	+	Mix
WD1005	TCC ATG AGC TTC CTG AGC TT	-	+
WD1006	TCC ATG ACG TTC CTG ACG TT	+	-
WD1007	ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG	+	+
	TCG TCG TTT TGT CGT TTT GTC GTT	+	+

20 * alternatively referred to as WD001-WD007

In the above table a + in the Thio column indicates the presence of a thioate modification. 'Mix' indicates a mixture of thioate modification and sequence without thioate modification (the asterisks indicate the linkages with a thioate modification). A - in the Thio column indicates absence of a thioate modification. A + in the CpG column indicates a the presence of a CpG motif and a - in the CpG column indicates absence of a CpG motif. For example WD1005 contains a GpC rather than a CpG motif, thus it is marked with a - in the CpG column of the table. WD1007 contains a palindromic motif (GACGTC) as well as other non-palindromic CpG sequences. This is also within the scope of a CpG oligonucleotide as the term is used in the present application.

The oligonucleotides utilised in the present invention may be synthesized by any method known in the art (eg EP 0 468 520). Conveniently, such oligonucleotides may be synthesized utilising an automated synthesizer. Methods for producing phosphorothioate oligonucleotides or phosphorodithioate are described in US patent 5,666,153, US patent 5,278,302 and WO95/26204.

The amount of protein in each vaccine does is selected as an amount which induces an immunoprotective response without significant, adverse side effects in typical vaccinees. Such amount will vary depending upon which specific immunogen is employed and how it is presented. Generally, it is expected that each dose will comprise 1-1000 µg of protein, preferably 2-100 µg, most preferably 5-50 µg. An optimal amount for a particular vaccine can be ascertained by standard studies involving observation of appropriate immune responses in subjects. Following an initial vaccination, subjects may receive one or several booster immunisations adequately spaced.

It is also possible to pre-administer the CpG oligonucleotide as a 'priming formulation' shortly prior to vaccination with the malaria antigen, for example 1 day before.

Accordingly, according to another aspect of the invention, there is provided a method for the prevention or amelioration of plasmodium infection in a patient, comprising administering an effective amount of either a malaria antigen and a CpG oligonucleotide (as hereinabove defined) or an effective amount of the CpG
5 oligonucleotide followed after a suitable time by an effective amount of a malaria antigen.

There is also provided a kit comprising effective amounts of a CpG oligonucleotide-containing formulation for use as a priming formulation for pre-administration to
10 human patients and a malaria antigen for injection at some suitable time later, as described hereinabove.

Preferred CpG oligonucleotides are those indicated in the table hereinabove.

15 Suitably the CpG will be present in the range 10 µg per dose to 1000 µg, preferably 10-100µg, especially 25-75 µg, for example 50 µg per dose.

Suitably the vaccine used in the present invention may comprise a carrier such as an aluminium salt, eg aluminium hydroxide [Al(OH)₃], aluminium phosphate or
20 aluminium phosphate sulfate (alum), or a non-toxic oil in water emulsion or a mixture thereof.

If an aluminium salt (preferably aluminium hydroxide) is used as a carrier it is generally present in the range of 50 to 100 µg, preferably 100 to 500 µg per dose.
25

Non-toxic oil in water emulsions preferably contain a non-toxic oil, eg squalene and an emulsifier such as (polysorbitan monoleate) Tween 80, in an aqueous carrier such as phosphate buffered saline.

30 If desired the vaccine used in the present invention may comprise an additional adjuvant, preferably a saponin adjuvant such as QS21 as described for example in

WO 9517210, optionally in the presence of a sterol, such as cholesterol as described for example in PCT/EP96/01464. The vaccine of the invention may also comprise monophosphoryl lipid A and derivatives thereof known in the art. A preferred derivative is 3 de-O-acylated monophosphoryl lipid A, described in British Patent
5 No. 2220211.

Accordingly vaccine formulations of the present invention may additionally comprise other pharmaceutical excipients or immunostimulants. In a preferred embodiment the vaccine formulation additionally comprises an aluminium salt,
10 preferably aluminium hydroxide.

The present invention will now be described with reference to the following examples:

15 **IMMUNOGENICITY STUDIES USING RTS,S FORMULATED WITH CPG OR CPG/ALUM**

Evaluation of CpG and CpG/alum in mice

Experiment outline

20 An immunogenicity study was conducted to evaluate the ability of CpG to serve as an adjuvant for cytotoxic T lymphocyte (CTL) induction. Groups of mice were immunized with RTS,S formulated with CpG oligonucleotide alone or in combination with aluminum hydroxide. After two immunizations spleen cells were examined for the presence of HbsAg-specific effector cells.

25

Table 1: Groups of mice

Group	antigen	adjuvant
1	RTS,S	CpG/alum
2	RTS,S	CpG

Formulation*Component batches used:*

COMPONENT	BRAND	BATCH NUMBER	CONCENTRATION (MG/ML)	BUFFER
RTSS		24R51	0.664	P/N 6.8
Al(OH) ₃	Superfos	97A0027	10.380	H ₂ O
CpG (WD1001)	Eurogen etech	324255R1	5	H ₂ O

5 Formulation process:

Formulations were prepared 3 days before each injection. All incubations were carried out at room temperature with agitation.

CpG/alum group 1 (500µl/dose)

- 10 RTS, S (8.7µg) and gp120 (8.7µg) were adsorbed on 100µg of Al(OH)₃ or AlPO₄ for 1 hour. The formulation was buffered with a 10-fold concentrated PO₄/NaCl pH 6.8 solution before addition of 100µg of CpG (WD1001). After 15 min, 50 µg/ml of thiomersal was added as preservative.

15 CpG group 2 (500µl/dose)

RTS,S (8.7µg) and Gp120(8.7µg) were diluted in PBS pH 6.8 before addition of 100µg of CpG (WD1001). After 5 min, 50 µg/ml of thiomersal was added as preservative.

20 Immunological methods

Nine Balb/C mice per group received into the hind footpads 100 µl vaccine twice at a two-week-interval. Two weeks later spleen cells were harvested and used to determine the induction of HBsAg-specific CTL.

For CTL analysis cells were cultured for 7 days in 6-well plates in the presence of 10 µg per ml of synthetic peptide pCMI003 corresponding to an HBsAg CTL epitope (Schirmbeck et al., 1995). At the end of the culture period cells were assessed in duplicate for HBsAg-specific cytolytic activity in standard [⁵¹Cr]-release assays using control and S-transfected P815 cells. Minimum and maximum release were determined with target cells without effector cells and by the addition of 3 % (v/v) Triton X-100, respectively. Results are expressed as % [⁵¹Cr]-release (cpm of exp. culture - cpm of spont. release / cpm of max. release - cpm of spont. release).

10 Results

Spleen cells from both groups of mice exhibited HBsAg-specific effector cell function in ⁵¹Cr release assays (Figure 1). Untransfected P815 target cells were lysed to a much lesser degree than similar target cells expressing the HBV s gene. Lysis of the target cells diminished with decreasing effector to target cell ratios.

15

Conclusions

Immunization with RTS,S in combination with CpG or CpG/alum induces HBsAg-specific CTL in mice.

20 EVALUATION OF CPG AND CPG/ALUM IN RHESUS MONKEYS

Experiment outline

An immunogenicity study was conducted to evaluate the adjuvant effect of CpG in non-human primates. Groups of five monkeys were immunized twice with RTS,S in combination with CpG or CpG/alum. After the second immunization the immune response of the animals was assessed. Antibodies to HBsAg and lymphoproliferative as well as cytokine responses were evaluated.

30

Table 2. Groups of monkeys

Group	antigen	adjuvant
1	RTS,S	CpG/alum
2	RTS,S	CpG

Formulation*Component batches used:*

5

COMPONENT	BRAND	BATCH NUMBER	CONCENTRATION (MG/ML)	Buffer
RTS,S		ERTS1X058	1.372	P/N 6.8
Al(OH) ₃	Superfos	96A0089	10.380	H ₂ O
CpG		WD1001	5	H ₂ O

Formulation process:

Formulations were prepared one day before each injection. All incubations were carried out at room temperature with agitation.

10 *CpG/alum group 1 (500µl/dose)*

RTS,S (50µg) was adsorbed on 500µg of Al(OH)₃ for 1 hour. The formulation was buffered with a 10-fold concentrated PO₄/NaCl pH 6.8 solution before addition of 500µg of CpG (WD1001). After 15 min, 50 µg/ml of thiomersal was added as preservative.

15

H₂O + Al(OH)₃ + RTS,S-1H-10xPN-15m-CpG-15m-Thio

CpG group 2 (500µl/dose)

RTS,S (50µg) was diluted in PO₄/NaCl buffer pH 6.8 before addition of 500µg of CpG WD1001. After 15 min, 50 µg/ml of thiomersal was added as preservative.

20

H₂O + RTSS + 10xPN-15m-CpG-15m-Thio

Immunological methods

Five rhesus monkeys (*Macaca mulatta*) per group were immunized twice intramuscularly with 500 μ l of vaccine at a four-week-interval. Sera and peripheral
5 blood mononuclear cells (PBMC) were taken at several occasions.

HBsAg-specific antibodies in monkey sera were determined in a radio immuno assay (RIA, Abbott) according to the manufacturer's instructions.

10 Lymphoproliferation was assessed by using density gradient-purified PBMC from immunized rhesus monkeys. Cells were seeded in quadruplicates at 1×10^5 in 100 μ l RPMI/5 % FCS per well in round bottom 96 well plates. Then another 100 μ l of medium alone or containing soluble RTS,S (10 μ g/ml) were added and parallel
15 cultures were incubated for 48 hrs. Thereafter, 100 μ l culture supernatant were replaced by fresh medium containing 1 μ Ci [3 H]-thymidine. After 16 hrs cells were harvested onto filter plates and incorporated radioactivity was determined in a β -counter. Results are expressed in cpm and in stimulation indices (SI, = cpm
antigen-containing cultures/cpm medium alone cultures), SI greater than 3 are considered as a positive response.

20 Flat bottom 96 well plates were prepared by coating an IFN- γ -specific capture antibody in 50 μ l PBS for 4 hrs at 37 °C. The plates were washed three times and PBMC were seeded similar to lymphoproliferation assays. After 48 hrs of culture the plates were washed thrice with PBS/0.05 % Tween 20 and 50 μ l of biotinylated
25 secondary IFN- γ -specific antibody diluted in PBS/Tween/1 % FCS were added for 2 hrs. The plates were washed again and a gold-conjugated α -biotin antibody was incubated for 1 hr. After additional washings the ELIspots were visualized by using a silver enhancing kit (50 μ l per well). The reaction was stopped after approx. 30 min by adding deionized water. Cytokine-secreting cells were enumerated by
30 microscopic examination.

Results

Analysis of HBsAg-specific antibodies in sera of the monkeys revealed that all animals in the two groups had developed specific immune responses (Figure 2). Some responses were detectable already after one immunization. Interestingly, these responses were boosted by the second immunization only in group 1, while titers in group 2 remained more or less constant.

Induction of specific lymphoproliferation by immunization with RTS,S in combination with CpG or CpG/alum was evaluated before immunization and 6 days post secondary immunization. All 10 animals did not exhibit any specific lymphoproliferation ($SI > 3$) at the study start (data not shown). In contrast, all animals in group 1 possessed strong lymphoproliferative responses 6 days post boost immunization (Figure 3). All animals from group 2 did, however, remain negative in this analysis.

15

The presence of RTS,S-specific IFN- γ -secreting cells was investigated in all monkeys before immunization and 6 days after the second dose. IFN- γ -secreting cells could not be evaluated from pre-immunization samples due to technical difficulties. However, such cells were detectable after secondary immunization (Figure 4). All animals in group 1 exhibited a positive response, while only one animal in group 2 was positive.

20

Conclusions

Immunization with RTS,S in combination with CpG induces immune responses in non-human primates. After two immunizations CpG alone induces low level HB sAg-specific antibodies, while CpG combined with alum induces high titer antibodies as well as vigorous lymphoproliferative and IFN- γ responses.

25

Figure legends

Figure 1: CTL activity of spleen cells from immunized mice. Effector cell activity was assessed by examining ^{51}Cr release of P815 cells (open circles) or *s*-transfected P815 cells (closed circles).

5

Figure 2: HBsAg-specific antibody responses in immunized rhesus monkeys. Specific antibodies were evaluated using a commercially available RIA. Individual values from multiple time points for each animal are shown in the table, and group averages are shown in the table and as a graphic.

10

Figure 3: RTS,S-specific lymphoproliferation in immunized rhesus monkeys 6 days post second immunization. PBMC were stimulated with RTS,S antigen and lymphoproliferative responses were measured by ^3H -thymidine incorporation. Results are expressed in cpm and as SI.

15

Figure 4: RTS,S-specific IFN- γ -secreting cells from immunized rhesus monkeys. IFN- γ -secreting cells were visualized by the ELISpot method. Cytokine-secreting cells resulting in a colored spot were enumerated by microscopic examination and results are expressed semi-quantitatively (- = 0-5, + = 5-15, ++ = 15-35, +++ = 35-50, ++++ = > 50).

20

References

Schirmbeck, R., Boehm, W., Melber, K., Reimann, J. (1995). Processing of exogenous heat-aggregated (denatured) and particulate (native) Hepatitis B surface antigen for class I-restricted epitope presentation. J. Immunol. 155:4676-4684.

25

Claims:


1. A vaccine formulation comprising a malaria antigen and an immunostimulatory
5 CpG oligonucleotide.
2. A vaccine as claimed in claim 1 wherein the antigen is selected from the group,
RTS, RTS*, TRAP or immunologically equivalent derivatives thereof.
3. A vaccine as claimed in claim 1 or 2 wherein the vaccine comprises TRAP or
immunologically equivalent derivative and one of RTS or RTS*.
- 10 4. A vaccine formulation as claimed herein additionally comprising an aluminium
salt, 3 de-O-acylated monophosphoryl lipid A or a saponin adjuvant.
5. A vaccine as claimed herein wherein the oligonucleotide comprises two CpG
dinucleotides.
6. A vaccine as claimed herein wherein the CpG oligonucleotide is between 15-45
15 nucleotides in length.
7. A vaccine as claimed herein wherein the CpG oligonucleotide comprises at least
one phosphorothioate internucleotide bond.
8. A vaccine as claimed herein wherein the oligonucleotide is selected from the
group:

20

Oligo (internal designation)	5'-SEQUENCE-3'	CpG	Thio
WD1001	TCC ATG ACG TTC CTG ACG TT	+	+
WD1002	TCT CCC AGC GTG CGC CAT	+	+
WD1003	ACC GAT AAC GTT GCC GGT GAC G	+	-
WD1004	G*G*G GTC AAC GTT GAG* G*G*G* G*G	+	Mix
WD1005	TCC ATG AGC TTC CTG AGC TT	-	+
WD1006	TCC ATG ACG TTC CTG ACG TT	+	-
WD1007	ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG	+	+
	TCG TCG TTT TGT CGT TTT GTC GTT	+	+

9. A method for the prevention or amelioration of plasmodium infection in a patient, comprising administering an effective amount of a vaccine of any one of claims 1 to 8 to a patient.
- 5 10. A vaccine as claimed herein for use as a medicament.
11. A method of producing a vaccine as claimed in any one of claims 1 to 8 comprising admixing a malarial antigen and a CpG immunostimulatory oligonucleotide.
- 10
12. A method for the prevention or amelioration of plasmodium infection in a patient, comprising administering an effective amount of a CpG oligonucleotide followed after a suitable time by an effective amount of a malaria antigen.


1 / 4

**Fig 1:**

2/4

Fig 2:**Titer anti-HBs (m UI/ml-RIA)**

		d -12	average	14 d post I	average	7 d post II	average	28 d post II	average
Group 1	N5V	<1		115		921		1352	
	I055	<1		76		6424		1112	
	R441	<1	8	222	193	369	1907	431	888
	BXJ	8		121		862		546	
	CPJ	<1		430		960		997	
Group 2	NJ5	<1		<1		17		11	
	X006	4		4		51		37	
	I041	<1	4	11	29	42	71	5	35
	R361	<1		2		68		62	
	BBE	3		100		175		61	

Fig. 2a**Fig. 2b**

		-12	0	14	28	34	56
Group 1		8	193	1907	888		
Group 2		4	29	71	35		

Fig. 2c

Fig 3:




Fig. 3a




Fig. 3c




Fig. 3b




Fig. 3d

Fig 4:

Group 1						
	N5V	I055	R441	BXJ	CPJ	
RTS,S	+	++	++	++	++	5/5
Unstimulated cells	-	-	-	-	-	

Group 2						
	NJ5	X006	I041	R361	BBE	
RTS,S	-	+	+++	+	-	1/5
Unstimulated cells	-	+	+	+	-	

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number
WO 01/00231 A3

(51) International Patent Classification⁷: A61K 39/39, A61P 33/06 // (A61K 39/39, 39:005)

(21) International Application Number: PCT/EP00/05841

(22) International Filing Date: 23 June 2000 (23.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9915204.3 29 June 1999 (29.06.1999) GB

(71) Applicant (*for all designated States except US*):
SMITHKLINE BEECHAM BIOLOGICALS S.A.
[BE/BE]; Rue de L'Institut 89, B-1330 Rixensart (BE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): COHEN, Joseph
[US/BE]; SmithKline Beecham Biologicals S.A., Rue de
l'Institut 89, B-1330 Rixensart (BE). GARCON, Nathalie
[FR/BE]; SmithKline Beecham Biologicals S.A., Rue
de l'Institut 89, B-1330 Rixensart (BE). VOSS, Gerald
[DE/BE]; SmithKline Beecham Biologicals S.A., Rue de
l'Institut 89, B-1330 Rixensart (BE).

(74) Agent: TYRRELL, Arthur, William, Russell; SmithK-
line Beecham, Corporate Intellectual Property, Two New
Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

(88) Date of publication of the international search report:
5 July 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF CPG AS AN ADJUVANT FOR MALARIA VACCINE

(57) Abstract: A vaccine formulation for the prevention or amelioration of plasmodium infection in humans is provided. The vaccine comprises a malaria antigen, especially a protein which comprises a portion of the CS protein of *P. falciparum* fused in frame via a linear linker to the N-terminal of HBsAg, and an immunostimulatory CpG oligonucleotide. Methods for making the vaccine formulation of the invention are described. Patients may also be treated by pre-administration of the CpG oligonucleotide prior to administration of the malaria antigen.

WO 01/00231 A3

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/05841

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K39/39 A61P33/06 //(A61K39/39, 39:005)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, MEDLINE, SCISEARCH, EMBASE, CHEM ABS Data, WPI Data, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 05355 A (SMITHKLINE BEECHAM BIOLOG ; COHEN JOSEPH (BE)) 12 February 1998 (1998-02-12) cited in the application the whole document	1-12
Y	WO 99 11241 A (SMITHKLINE BEECHAM BIOLOG ; GARCON NATHALIE (BE); MOMIN PATRICIA MA) 11 March 1999 (1999-03-11) the whole document	1-12
Y	WO 96 02555 A (UNIV IOWA RES FOUND) 1 February 1996 (1996-02-01) cited in the application abstract claims 1,5,10,18	1-12

	---/---	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

8 document member of the same patent family

Date of the actual completion of the international search

19 January 2001

Date of mailing of the international search report

26/01/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Niemann, F

INTERNATIONAL SEARCH REPORT

International Publication No.

PCT/EP 00/05841

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 40100 A (DAVIS HEATHER L ; OTTAWA CIVIC LOEB RESEARCH INS (CA); QIAGEN GMBH) 17 September 1998 (1998-09-17) abstract; claim 16 ---	8
A	WO 98 37919 A (UNIV IOWA RES FOUND) 3 September 1998 (1998-09-03) abstract page 27, line 20 ---	8
P,X	WO 00 23105 A (SMITHKLINE BEECHAM BIOLOG ; GARCON NATHALIE (BE)) 27 April 2000 (2000-04-27) abstract; claims 1-26 ---	1-7,9-12
E	WO 00 62800 A (SMITHKLINE BEECHAM BIOLOG ; FRIEDE MARTIN (BE); GARCON NATHALIE (BE)) 26 October 2000 (2000-10-26) abstract page 15, line 12 - line 30 claims 1-10 ---	1-12
P,X	JONES T R ET AL: "Synthetic oligodeoxynucleotides containing CpG motifs enhance immunogenicity of a peptide malaria vaccine in Aotus monkeys" VACCINE, GB, BUTTERWORTH SCIENTIFIC. GUILDFORD, vol. 17, no. 23-24, 6 August 1999 (1999-08-06), pages 3065-3071, XP004173617 ISSN: 0264-410X the whole document -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter Application No

PCT/EP 00/05841

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9805355 A	12-02-1998	AU 706303 B	10-06-1999
		AU 4204097 A	25-02-1998
		BG 103141 A	31-08-1999
		BR 9710913 A	17-08-1999
		CN 1231613 A	13-10-1999
		CZ 9900292 A	16-06-1999
		EP 0957933 A	24-11-1999
		NO 990464 A	01-02-1999
		PL 331424 A	19-07-1999
		SK 11599 A	13-03-2000
WO 9911241 A	11-03-1999	AU 1145699 A	22-03-1999
		EP 1009382 A	21-06-2000
WO 9602555 A	01-02-1996	AU 713040 B	18-11-1999
		AU 1912795 A	16-02-1996
		CA 2194761 A	01-02-1996
		EP 0772619 A	14-05-1997
		JP 10506265 T	23-06-1998
		US 6008200 A	28-12-1999
WO 9840100 A	17-09-1998	AU 6759598 A	29-09-1998
		EP 1005368 A	07-06-2000
WO 9837919 A	03-09-1998	AU 6667498 A	18-09-1998
		EP 1039935 A	04-10-2000
WO 0023105 A	27-04-2000	AU 1151800 A	08-05-2000
WO 0062800 A	26-10-2000	NONE	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)